



This document contains information that will assist the owners of a small egg laying flocks in ensuring that locally produced eggs are of high quality and are safe for Virginia's consumers.

Coop and Nest Management

Keep the laying flock in a secure fenced area separated from wild birds that can transmit disease. Provide them nest boxes so they cannot hide their eggs or nest anywhere they choose. If hens are allowed to nest wherever they choose, you will not know how old eggs are when you find them or what they have been in contact with — if you can find them at all. Eggs laid in nest boxes will also be cleaner.

Keep birds safe. Don't attract wildlife to your flock by allowing birds to range free or by leaving feed or waste materials out in the open. Bears, dogs, foxes and birds of prey all love domestic poultry. Consider completely enclosing your coop and runs to prevent these predators from preying on your flock. This can be as easy as putting netting over the fenced yard. Once predators discover an easy meal, they will be back causing ongoing problems for you and your birds.

Provide a clean environment. Keeping the flock's environment clean and dry will help keep your eggs clean. A muddy outside run, dirty or damp litter and dirty nesting material will result in dirty, stained eggs. Clean out the nest boxes and add deep, clean litter at least every two weeks. Clean out wet litter in the coop and make sure the outside run area has good drainage and is not overgrazed.

Provide adequate nest space. Supply a minimum of four nesting boxes for flocks of 15 hens or fewer. For larger flocks provide one nest for every four to five hens in the flock. This will help limit egg breakage from normal traffic and daily egg laying. Make sure nests have a deep, clean layer of litter to prevent breakage and help absorb waste or broken egg material. Place nest boxes in dimly lit areas to help reduce or prevent egg pecking.

Make sure you have perches. Part of the bird's natural behavior is to roost. Providing flat wooden perches at least 4 inches across will allow the birds a feeling of security and comfort. Perches keep the birds feet warmer and help eliminate health problems like bumble foot.

Collecting the Eggs

A hen's laying time will vary depending on the temperature of the coop and at what time the hens are exposed to light. It is best to collect the eggs as soon as possible after they are laid, so get to know your flock's cycle. The longer the egg is allowed to stay in the nest, the more likely it is that the egg will get dirty, broken or lose interior quality. Collecting eggs at least twice daily is advisable.

A Few Tips for Layer House Management

Rotate range areas often or allow enough area for birds in outside runs to prevent large dirt and mud areas from forming by overgrazing. Electric poultry fencing or chicken tractors (portable coop and run structures) can be helpful in rotating areas.

Provide adequate lighting and a heat source during cold weather. Hens need a minimum of 15 hours of light to continue laying through the dark months. It's best to add light in the morning and not the evening in order to help maintain egg production and to make sure the birds aren't suddenly caught out in the dark when the lights switch off. Provide enough light so that newsprint can be read anywhere in the coop.

Provide hens with readily available grit and oyster shells to help strengthen the egg shells.

Store feed in a metal container and keep feeders up off the ground to discourage rodents.

Proper Egg Cleaning and Handling

Collect eggs and brush off any debris adhering to the egg shell, and then put them in an easy-to-clean container, such as a coated wire basket or plastic egg flat. This will prevent stains from rusted metal and contamination from other materials that are difficult to clean and disinfect.

Do not stack eggs too high. If collecting in baskets, do not stack eggs more than five layers deep. If using plastic flats, do not stack more than six flats. If you stack eggs too deep you will increase breakage.

Never cool eggs rapidly before they are cleaned. The eggshell will contract and pull any dirt or bacteria on the surface deep into the pores as the egg cools. Try to keep the temperature relatively constant until they are washed.

Wash eggs as soon as you collect them. This helps limit contamination and loss of interior quality.

Wash eggs with water that is at least 90°F. This will make the egg contents swell and push the dirt away from the pores of the egg. If you have extremely dirty eggs, a mild detergent approved by the FDA for washing eggs can be used. Never let eggs soak in water. Soaking eggs in water for as little as one to three minutes can allow bacteria to penetrate the shell and contaminate the egg. The eggs can be sanitized to further reduce the bacteria numbers on the shell by using one tablespoon of unscented household chlorine bleach per gallon of water.


Cool and dry eggs quickly after washing. Store eggs, large end up, at 45°F and at 75 percent relative humidity. If eggs sit at room temperature (75°F), they can drop as much as one grade per day. If fertile eggs are kept at a temperature above 85°F for more than a few hours, the germinal disc (embryo) can start to develop. If fertile eggs are kept above 85°F over two days, the blood vessels of the embryo may become visible. Eggs stored properly in their own carton or other stable environment will maintain a quality of Grade A for at least four weeks.

Sorting and Grading Eggs

It is best that you sort the eggs before you store, sell or consume them. The easiest way to sort eggs is to candle them with a bright light. This process can help you eliminate cracked eggs, eggs with foreign matter inside (such as blood spots), or other poor quality eggs.

Candling eggs

Candling lights are commercially available, but candling can be done with less specialized equipment. A high intensity flashlight can be used with practice (a modern LED flashlight may be a good choice, since light output can be adequate and the light produces little heat). Working in a darkened room for optimal results, hold the egg up to the candling light in a slanting position. You will see the air cell, the yolk and the white; the air cell is almost always in the large end of the egg. Put the large end next to the candling light. Hold the egg between your thumb and first two fingers. By turning your wrist quickly, you can cause the inside of the egg to whirl. This will tell you a great deal about the yolk and white. When you are learning to candle, you will find it helpful to break and observe any eggs you are in doubt about.



USDA grade standard: Use the specifications given in the table to the right to determine the grade of an egg. Consider shell, air cell depth, yolk outline and albumen quality.

Shell: A normal, high-quality egg has an oval shape with one end larger than the other. The shell should be sound with no cracks, apparent thin spots or ridges that affect the soundness of the shell.

Identifying cracks: Cracked eggs will appear to have a white line somewhere on the shell. These cracks will open if you apply slight pressure to the shell. Remove cracked eggs and consume them as soon as possible or discard.

Air cell depth: The depth of the air cell is the distance from its top to its bottom when the egg is held with the air cell up. In a fresh egg, the air cell is small, not more than 1/8 inch deep. As the egg ages, evaporation takes place and the air cell becomes larger, resulting in a lower grade egg.

White, or albumen: The character or condition of the white, or albumen, can be determined by the behavior of the yolk when the egg is candled. If the yolk retains its position in the center when the egg is twirled, the white is usually firm and thick.

Yolk: The yolk of a fresh, high-quality egg will be surrounded by a rather dense layer of albumen, or white. Therefore, it moves only slightly away from the center of the egg when it is twirled in front of the candling light, and the yolk outline is only slightly defined. As the albumen thins, the yolk tends to move more freely and closer to the shell and the defined outline is more easily seen. A more visible yolk when candled indicates a lower quality egg.

Spots: Eggs with blood or meat spots more than 1/8 inch in diameter are classified as inedible. Eggs with small spots collectively less than 1/8 inch in diameter should be classified as Grade B. The *chalaza* (whitish, ropelike protein on each side of the yolk that suspends it in the albumen) is distinguished from a meat spot by a bright area of refracted light that accompanies its darker shadow. Blood spot eggs can be consumed without harm; however, people may find the appearance undesirable.



Check this out!
↳

TABLE 2. Summary of U.S. Standards for Quality of Individual Shell Eggs.

SPECIFICATIONS FOR EACH QUALITY FACTOR			
Quality Factor	AA Quality	A Quality	B Quality
Shell	Clean Unbroken. Practically Normal.	Clean Unbroken. Practically Normal.	Clean to slightly stained.* Unbroken. Abnormal.
Air Cell	1/8 inch or less in depth. Unlimited movement and free or bubbly.	3/16 inch or less in depth. Unlimited movement and free or bubbly.	Over 3/16 inch in depth. Unlimited movement and free or bubbly.
White	Clear. Firm.	Clear. Reasonably firm.	Weak and watery. Small blood and meat spots present.**
Yolk	Outline slightly defined. Practically free from defects.	Outline fairly well defined. Practically free from defects.	Outline plainly visible. Enlarged and flattened. Clearly visible germ development but no blood. Other serious defects.
For eggs with dirty or broken shells, the standards of quality provide two additional qualities.			
	Dirty	Check	
	Unbroken. Adhering dirt or foreign material, prominent stains, moderate stained areas in excess of B quality.	Broken or cracked shell, but membranes intact, not leaking.***	
	* Moderately stained areas permitted (1/32 of surface if localized, or 1/16 if scattered.) ** If they are small (aggregating not more than 1/8 inch in diameter). *** Leaker has broken or cracked shell membranes, and contents leaking or free to leak.		

This table may be found in the USDA "Egg Grading Manual". Please see the manual for more in-depth grading and egg information.

GRADING EGGS MAY BE DONE BY THE PRODUCER
One does not need to be a USDA grader to grade eggs!



Sizing eggs: Eggs can be sized for marketing since consumers often prefer medium, large and extra large eggs. Weigh your eggs and use the specifications in the table below to determine size classifications. Egg scales can be purchased through a poultry supply company, but any accurate scale will work. Sizing your eggs is optional.

Egg Sizing Chart	
Size or weight class	Minimum net weight per dozen
Jumbo	30 ounces
Extra large	27 ounces
Large	24 ounces
Medium	21 ounces
Small	18 ounces
Peewee	15 ounces

Maintaining quality

Refrigeration maintains the quality of the egg. A holding temperature of $\leq 45^{\circ}\text{F}$ will slow the deterioration of the egg quality and reduce the growth of bacterial pathogens such as *Salmonella*.

The egg is laid at 105°F , the normal body temperature of the hen. As the egg cools, the albumen and yolk shrink away from the shell, forming the air cell. Over time, the albumen loses water and continues to shrink, causing the air cell to increase in size and, thus indicating a decline in quality.

Over time, the structure of the white and yolk will begin to break down, decreasing the quality of the egg and resulting in an egg with a thinner, more watery albumen, or egg white, and a softer, flatter and weaker egg yolk. If broken open, the egg will spread over a large area with a clear, watery white with reduced leavening and thickening powers; the yolk will be flatter and more easily broken. Using poor quality eggs in cooking or baking may cause a recipe to fail.

Storing eggs

Store eggs in an egg carton with the small end down to keep the air cell stable.

Date the carton so you can use or sell the oldest eggs first, and rotate your extra eggs. Try to use or sell all eggs before they are three weeks old.

Store eggs in a refrigerator with an ambient temperature of $\leq 45^{\circ}\text{F}$ and 70-85% humidity. Place a thermometer in the refrigerator to track the temperature of your fridge. Never hold eggs at or above room temperature or at low humidity longer than necessary. Leaving eggs in a warm, dry environment will cause quality to drop quickly.

Never store eggs with materials that have an odor. Eggs will pick up the odors of apples, fish, onions, potatoes and other food or chemicals with distinct odors.

Package (carton) labels

If the eggs are packaged to be sold at a food establishment, they must meet food labeling laws and regulations. To be in compliance, **your eggs should be labeled with the following information in English**, using at least a ¼-inch font size:

- Statement of identity (eggs)
- Grade and size/weight class
- Numeric count (for example, one dozen, two dozen)
- Name and address of the producer (farm name) distributor, packer or retailer. If the name placed on the label can be found in a telephone directory, only the city, state and ZIP code need to be included. Otherwise, the street address must be included.
- The words "Keep Refrigerated"
- Safe handling instructions, with the statement:
- SAFE HANDLING INSTRUCTIONS:** To prevent illness from bacteria: keep eggs refrigerated, cook eggs until yolks are firm and cook foods containing eggs thoroughly.

ONE DOZEN	Good Egg Farm	SAFE HANDLING INSTRUCTIONS: To prevent illness from
GRADE A EGGS	1234 Amber Lane	bacteria: keep eggs refrigerated, cook eggs until yolks
KEEP REFRIGERATED	Farmland, VA 12345	are firm and cook foods containing eggs thoroughly.

Example of label to be secure to top of egg carton.

It is acceptable to recycle cartons if the cartons are clean and in good condition and if all labeling information that does not apply is removed. Egg cartons must be free from inspection or labeling marks from previous use.

Specialty Labels



Organic: Eggs **cannot** be labeled or represented as organic unless the operation has been certified by an accredited National Organic Program certifier. <http://www.ams.usda.gov/AMSv1.0/nop>



Virginia's Finest: Eggs can be sold with the Virginia's Finest on their label **ONLY** if they are enrolled in the Virginia's Finest Program. <http://www.vafinest.com/>

Transporting and displaying eggs

Eggs should be kept at a temperature $\leq 45^{\circ}\text{F}$ during transportation to the point of sale. This can be done by keeping the eggs under refrigeration until transport, then using a cooler with gel packs to keep the eggs cool. If the eggs are to be sold at a farmers market or roadside stand, use mechanical refrigeration, such as a small refrigerator or plug-in cooler, if possible. A thermometer should be placed in the cooler so that the temperature can be monitored to ensure that the air temperature in the cooler is $\leq 45^{\circ}\text{F}$.

Selling eggs to a food establishment

If the eggs are delivered to or sold to someone who is not the end consumer, such as a food establishment, they must be received at the establishment in refrigerated equipment that maintains an ambient air temperature of 45°F degrees or below. The eggs must be clean and sound and not exceed the restricted egg tolerances for the U.S. consumer Grade B (see egg grading section). The eggs must be properly labeled (see section on package labels).

References

Alaska Cooperative Extension Service, *Safe Egg Handling for Small Egg Laying Flocks and Operations*:
<http://www.uaf.edu/files/ces/publications-db/catalog/anr/LPM-00344.pdf>

USDA Agriculture Marketing Service, National Organic Program:
<http://www.ams.usda.gov/AMSV1.0/nop>

USDA Agriculture Marketing Service, Egg Grading Manual:
<http://www.ams.usda.gov/AMSV1.0/getfile?dDocName=STELPRDC5110016>

Virginia Cooperative Extension Service, *Proper Handling of Eggs: From Hen to Consumption*:
https://pubs.ext.vt.edu/2902/2902-1091/2902-1091_pdf.pdf

Virginia Department of Agriculture and Consumer Services Virginia's Finest Program:
<http://www.vafinest.com/>

Virginia Laws and Regulations:
<https://leg1.state.va.us/lis.htm>

Virginia's Finest Program
<http://www.vafinest.com/>